



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2020

Visual snow syndrome: a review on diagnosis, pathophysiology, and treatment

Traber, Ghislaine L ; Piccirelli, Marco ; Michels, Lars

Abstract: **PURPOSE OF REVIEW** Visual snow is considered a disorder of central visual processing resulting in a perturbed perception of constant bilateral whole-visual field flickering or pixelation. When associated with additional visual symptoms, it is referred to as visual snow syndrome. Its pathophysiology remains elusive. This review highlights the visual snow literature focusing on recent clinical studies that add to our understanding of its clinical picture, pathophysiology, and treatment. **RECENT FINDINGS** Clinical characterization of visual snow syndrome is evolving, including a suggested modification of diagnostic criteria. Regarding pathophysiology, two recent studies tested the hypothesis of dysfunctional visual processing and occipital cortex hyperexcitability using electrophysiology. Likewise, advanced functional imaging shows promise to allow further insights into disease mechanisms. A retrospective study now provides Class IV evidence for a possible benefit of lamotrigine in a minority of patients. **SUMMARY** Scientific understanding of visual snow syndrome is growing. Major challenges remain the subjective nature of the disease, its overlap with migraine, and the lack of quantifiable outcome measures, which are necessary for clinical trials. In that context, refined perceptual assessment, objective electrophysiological parameters, as well as advanced functional brain imaging studies, are promising tools in the pipeline.

DOI: <https://doi.org/10.1097/WCO.0000000000000768>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-177673>

Journal Article

Published Version

Originally published at:

Traber, Ghislaine L; Piccirelli, Marco; Michels, Lars (2020). Visual snow syndrome: a review on diagnosis, pathophysiology, and treatment. *Current Opinion in Neurology*, 33(1):74-78.

DOI: <https://doi.org/10.1097/WCO.0000000000000768>



Visual snow syndrome: a review on diagnosis, pathophysiology, and treatment

Ghislaine L. Traber^{a,b}, Marco Piccirelli^c, and Lars Michels^c

Purpose of review

Visual snow is considered a disorder of central visual processing resulting in a perturbed perception of constant bilateral whole-visual field flickering or pixelation. When associated with additional visual symptoms, it is referred to as *visual snow syndrome*. Its pathophysiology remains elusive. This review highlights the visual snow literature focusing on recent clinical studies that add to our understanding of its clinical picture, pathophysiology, and treatment.

Recent findings

Clinical characterization of visual snow syndrome is evolving, including a suggested modification of diagnostic criteria. Regarding pathophysiology, two recent studies tested the hypothesis of dysfunctional visual processing and occipital cortex hyperexcitability using electrophysiology. Likewise, advanced functional imaging shows promise to allow further insights into disease mechanisms. A retrospective study now provides Class IV evidence for a possible benefit of lamotrigine in a minority of patients.

Summary

Scientific understanding of visual snow syndrome is growing. Major challenges remain the subjective nature of the disease, its overlap with migraine, and the lack of quantifiable outcome measures, which are necessary for clinical trials. In that context, refined perceptual assessment, objective electrophysiological parameters, as well as advanced functional brain imaging studies, are promising tools in the pipeline.

Keywords

migraine, palinopsia, persistent visual phenomena, visual snow, visual static

INTRODUCTION

Since the first description of visual snow in the literature in 1995 [1], visual snow syndrome has emerged as a distinct clinical entity [2^{••}]. The hallmark of disease is the perception of constant bilateral whole-visual field flickering or pixelation, affecting the patients' daily life. *Visual snow syndrome* is defined as visual snow with at least two additional visual symptoms as outlined below. Symptom onset ranges from childhood [3,4] to elderly patients [5], but is usually reported in young adulthood, equally affecting men and women [6,7[•]]. There is an ongoing debate as to whether visual snow syndrome should be considered as an organic disease or rather as a heightened perception of normal sensory phenomena [8]. However, the surprising stereotypy of patients describing their symptoms, even in the absence of previous knowledge about this condition, speaks against it being psychogenic in nature. Visual snow syndrome, by definition, is a benign condition and should not be associated with signs of a progressive ophthalmic or neurological disease. On rare occasions, however,

visual snow symptoms have also been described in patients with serious neurological diseases such as Creutzfeldt-Jakob disease [9] or glycine receptor antibody syndrome [10]. The latter again seems to underpin a probable organic basis and emphasizes that careful adherence to diagnostic criteria, particularly the exclusion of other ophthalmic or neurological disease, is essential.

Early studies have focused on clinical characterization of visual snow syndrome [11–14]. The pathophysiology, however, remains still unclear, which goes hand in hand with the lack of effective treatment strategies. This review follows up on a

^aDepartment of Ophthalmology, University Hospital Basel, University of Basel, ^bInstitute of Molecular and Clinical Ophthalmology Basel (IOB), Basel and ^cDepartment of Neuroradiology, Clinical Neuroscience Center, University Hospital Zurich, University of Zurich, Zurich, Switzerland

Correspondence to Ghislaine L. Traber, MD, Department of Ophthalmology, University Hospital Basel, Mittlere Strasse 91, 4031 Basel, Switzerland. Tel: +41 61 265 8686; e-mail: ghislaine.traber@usb.ch

Curr Opin Neurol 2019, 32:000–000

DOI:10.1097/WCO.0000000000000768

KEY POINTS

- Thorough clinical characterization of visual snow syndrome has led to its recognition as a clinical entity.
- The pathophysiology remains elusive; however, bilateral pan-visual field involvement, frequently associated palinopsia, abnormal suprathreshold processing of contrast and luminance, electrophysiological findings as well as functional brain imaging point towards dysfunctional central visual processing.
- Major challenges remain the purely subjective nature of visual snow syndrome, its overlap with migraine, and most importantly, the lack of quantifiable outcome measures, which are prerequisite to design prospective clinical trials.
- Based on a retrospective chart review, there is Class IV evidence that a minority of patients may profit from a trial of lamotrigine.
- The majority of patients copes well with reassurance about the benign nature of visual snow syndrome.

comprehensive review article about visual snow syndrome by Puledda *et al.* [2¹¹] published in February 2018 in this journal. Our article highlights the most recent literature on visual snow syndrome with a focus on the clinical studies that add to our understanding and management of the condition. The objective of this review is to condense our current appreciation of the clinical picture, of the pathophysiology and available treatments.

We did a systematic literature search on PubMed up to August 2019 for articles on visual snow with the keywords ‘visual snow’ ‘visual snow syndrome’, ‘visual static’, and ‘persistent positive visual phenomena’.

VISUAL SNOW SYNDROME

The clinical picture

Based on their first systematic characterization of visual snow patients, preliminary criteria for visual snow syndrome were elaborated by Schankin *et al.* [11] and later suggested as definite clinical criteria [2¹¹]. These consist of

- (1) Visual snow lasting longer than 3 months: perceived as dynamic, continuous, tiny dots in the entire visual field.
- (2) At least two additional visual symptoms of the following four categories:
 - (a) Palinopsia: afterimages or trailing of moving objects.

- (b) Enhanced entoptic phenomena (modification suggested by Metzler and Robertson [15¹]: ‘other frequent or persistent visual phenomena’): excessive floaters in both eyes, excessive blue field entoptic phenomena, self-light of the eye, spontaneous photopsias.
 - (c) Photophobia.
 - (d) Nyctalopia.
- (3) Symptoms are not consistent with typical migraine visual aura.
 - (4) Symptoms are not better explained by another disorder.

Metzler and Robertson [15¹] recently challenged these criteria, namely section B (ii) ‘enhanced entoptic phenomena’ based on the fact that they had difficulty applying this section in the clinic. Instead, they proposed B (ii) to be labelled ‘other frequent or persistent visual phenomena’ arguing that ‘entoptic phenomena’ are generally considered to be phenomena arising from structures within the eye. Apart from the fact that photopsias do not necessarily originate from within the eye [16], the proposed mild modification of the visual snow syndrome criteria would allow accounting for otherwise unclassifiable phenomena described by some patients with visual snow. The latter include ‘halos around light-sources’, ‘geometric and coloured images that distort vision’, ‘metamorphopsias’, ‘straight lines moving across the visual field’, ‘water running down a window’, and ‘coloured clouds when closing eyes’ [11].

Irrespective of this semantic debate, visual snow syndrome criteria base on purely subjective descriptions of visual symptoms. Ophthalmic and neurological examination is typically normal [11], including perimetry [5,12,13,17], optical coherence tomography [17], electroretinography [11–13,17] as well as – until recently – visual evoked potentials (VEP) [11–13,17].

On brain imaging, left occipital bending has been mentioned in 4/14 patients [18¹¹]. Two studies found this structural peculiarity to be more frequent in patients with major depressive disorders (35.3 vs. 12.3%) and with bipolar disorders (35.3 vs. 8.3%) compared with healthy controls. However, the clinical significance of this finding in patients with visual snow is unclear. Otherwise, structural MRI findings are typically reported normal [12,13].

Commonly associated nonvisual symptoms are comorbid migraine, tinnitus, impaired concentration, lethargy, anxiety, depression, balance disorder, and tremor [5,7¹,13,19¹¹]. Migraine clearly is the major confounding factor and thus challenges visual snow research. It is not only a very common

comorbid disease (up to 58% of visual snow patients) [13,14,19²²], but probably also shares common disease mechanisms with visual snow [11,14], which has been discussed in more detail by Puledda *et al.* [2²²].

Pathophysiology

Visual snow syndrome is generally considered to involve dysfunctional central visual processing [2²²,6,7²²,15²²,19²²]. However, its cause remains largely unknown.

Lauschke *et al.* [13] tested colourimetry in 12 visual snow patients and found the yellow-blue colour spectrum to provide relief of symptoms in 10/12 patients. Thus, they consider visual snow to be a disorder of central colour-dependent processing involving the magnocellular pathway and suggested thalamocortical dysrhythmia as a possible underlying mechanism. The latter disease concept has been mostly discussed in the context of neurogenic pain, tinnitus, Parkinson's disease, and depression [20–22]. While being an interesting hypothesis, studies actually measuring thalamocortical dysrhythmia in visual snow patients are regrettably still missing.

Chen *et al.* [23] mentioned cortical hyperexcitability as a possible pathophysiological mechanism based on a magnetoencephalography study involving six patients with 'persistent visual aura', two of which had visual snow.

More recently, the concept of cortical hyperexcitability was supported in a behavioural study by McKendrick *et al.* [24]. Consistent with elevated spontaneous neural firing in the primary visual cortex, they found abnormal suprathreshold processing of contrast (centre-surround contrast matching) and luminance (luminance increment detection in noise) in 16 visual snow patients compared with 18 controls. However, no differences were found for global form and motion tasks, which are attributed to extrastriate processing. If confirmed in larger studies, measurement of perceptual differences in visual snow may hold potential for diagnostic clinical tests and monitoring of treatment effects in the future.

Eren *et al.* [25²²] tested the hypothesis of dysfunctional visual processing using pattern-reversal VEP in 18 affected patients compared with 18 migraineurs and 18 healthy controls. On a group level, they found a trend for reduced N75-P100 amplitudes in visual snow patients and a mildly, but significantly increased N145 latency compared with healthy controls independent of comorbid migraine. The fusiform gyrus is the supposed origin of the late phase P100 [26] and is adjacent to the lingual gyrus. Arguing that late phase P100

represents the transition to N145, – which was prolonged in the study – the authors claim to possibly show the first electrophysiological correlate to the reported hypermetabolism of the lingual gyrus [14]. While the line of argument regarding localization seems rather long-winded, the findings of Eren *et al.* might still further support the idea of visual snow as a disorder involving extrastriate cortical areas [25²²]. Of note, latencies of N75 and P100 did not differ between visual snow and controls, making dysfunction of the early cortical visual processing unlikely. In addition, there was no statistically significant difference of N75-P100 amplitude habituation to repetitive VEP stimulation (six consecutive blocks of 75 responses) between groups.

Based on related previous case reports [27,28], Yildiz *et al.* [18²²] prospectively investigated occipital cortex hyperexcitability by measuring the habituation response to repetitive pattern-reversal VEP consisting of 10 blocks of 100 averaged responses. Ten visual snow patients with migraine (VS_m), seven visual snow patients without migraine (VS_{wom}), and 12 healthy controls were recruited. They measured the N75-P100 VEP amplitude ratios of the 10th/1st block of the right and left eye. Contrary to the results of Eren *et al.*, this study found a loss of the habituation response in VS_m and VS_{wom} compared with the expected habituation in healthy controls. There was no statistically significant difference between VS_m and VS_{wom}, meaning that presence or absence of migraine did not influence the habituation response. Lack of habituation has been considered a neurophysiological hallmark of interictal migraine. However, the latter has been put into question recently [29]. Unfortunately, the study by Yildiz *et al.* [18²²] did not include a pure migraine control group without visual snow. The authors also assessed transcranial magnetic stimulation phosphene detection and found a lower left phosphene threshold in patients with visual snow. In the post-hoc analysis, the latter difference only reached statistical significance in VS_m compared with healthy controls. In sum, Yildiz *et al.* propose loss of VEP habituation and decreased phosphene threshold as possible surrogate markers indicating occipital cortex hyperexcitability.

It is understood that objective electrophysiological parameters would be extremely useful to assay benefits of treatment in visual snow. At present, results of electrophysiological studies seem to be differing [11–13,17,18²²,25²²] which might, however, partly be related to different examination protocols, for example, when assessing habituation. Larger studies are awaited to elucidate the field.

Functional brain imaging studies seem most appropriate to find evidence for dysfunctional

central visual processing. Apart from the occasional occipital bending mentioned above, brain imaging studies usually do not show specific *structural* abnormalities [11–13]. Jager *et al.* [30] investigated cerebral water diffusion and perfusion on MRI and found no regional functional changes in his two patients with visual snow. Schankin *et al.* [14] investigated seventeen patients with visual snow using [(18) F]-2-fluoro-2-deoxy-D-glucose PET and found hypermetabolism of the right lingual gyrus and left anterior cerebellar lobe. This finding is consistent with a disorder allocated downstream of the primary visual cortex.

To further investigate abnormal central visual processing, we are currently analysing resting state functional MRI (fMRI) of patients with visual snow compared with healthy controls within, but also outside of the visual cortex. Preliminary data in ten patients with visual snow and ten healthy controls was presented at the 45th North-American Neuro-Ophthalmology Society Meeting in Las Vegas, USA, in March 2019 (<https://www.nanosweb.org/files/Annual%20Meeting/2019/FINAL%20NANOS%20Brochure%20low%20res.pdf>). Our results point towards functional hyperconnectivity in brain areas involved in form and motion processing, as well as in areas associated with attention and working memory. These findings suggest, that visual snow patients not only have a metabolically altered early visual cortex (lingual gyrus) [31], but also show functional abnormalities in brain areas associated with the accompanying symptoms of visual snow, such as palinopsia (unpublished data).

In an attempt to explain why visual snow is frequently associated with additional symptoms, Metzler and Robertson [15[•]] discussed the concept of stochastic resonance, a nonlinear phenomenon where additional noise can improve the signal-to-noise ratio, thus lowering the detection threshold of another weak stimulus. The latter has not only been shown for stimuli involving the same sensory modality (e.g., vision) [32], but also for stimuli involving different senses (e.g., vision and hearing) [33]. The authors concluded that in a model of stochastic resonance, visual snow might, for example, enhance tinnitus or vice versa.

Patient management and treatment

Sharing our knowledge on visual snow and reassuring the patient about the benign nature of the condition often provides instant relief to the patient. Nevertheless, a minority of patients may request empiric treatment. Until recently, pharmacologic treatment was only based on case reports, small case series, or expert opinion [2^{••},6,28,34].

Recently, van Dongen *et al.* [19^{••}] evaluated treatment options in a retrospective case series of 58 visual snow patients, out of which 47 patients had visual snow syndrome. Given the lack of evidence, 29 patients chose no drug treatment. None of the remaining treated patients experienced complete remission under medication. Lamotrigine (2 × 50 mg) was prescribed most frequently ($n=26$) with partial remission of symptoms in 5/26 (19.2%). However, adverse events were reported in half of the patients (13/26) patients, including allergic reactions ($n=3$) and excessive daytime sleepiness ($n=1$). Eventually, only 4/26 patients continued lamotrigine treatment. One patient improved under topiramate ($n=4$), but still stopped the medication because of side effects. Valproate ($n=7$), acetazolamide ($n=2$) and flunarizine ($n=1$) did not improve symptoms in this case series. Overall, evidence for pharmacological treatment in visual snow remains scarce (Class IV evidence). Thus, treatment should be limited to selected patients only.

Treatment of comorbid diseases like migraine or mood disorders is certainly worth considering depending on the severity. However, there is no evidence suggesting benefit with regard to visual snow symptoms.

Recently, real-time fMRI neurofeedback has been proposed as a potential intervention for several neurological and psychiatric disorders [35–41]. As there is hypermetabolism in the lingual gyrus and growing evidence of dysfunctional connectivity in the brain, single region real-time fMRI neurofeedback may be a promising intervention strategy to normalize brain function in the future.

CONCLUSION

Research about visual snow syndrome has gone through an early phase of thorough clinical characterization leading to its recognition as a clinical entity. Overall, the disease is most likely related to dysfunctional central visual processing, which is also supported by the most recent clinical studies. Our scientific understanding of visual snow syndrome has started to grow as we now put bits and pieces of the puzzle together. However, major challenges remain, such as the purely subjective nature of visual snow syndrome, its overlap with migraine, and most importantly the lack of quantifiable and reliable outcome measures, which are prerequisite to design prospective clinical trials. In that context, measurement of refined perceptual differences, objective electrophysiological parameters, as well as advanced functional brain imaging studies including real-time fMRI neurofeedback, may hold

potential but need further investigations on a larger scale. In the meantime, we are left with Class IV evidence that a minority of patients may profit from a trial of oral lamotrigine and with the confidence that most patients cope very well with reassurance.

Acknowledgements

We thank Konrad P. Weber, MD, University Hospital Zurich, Switzerland, for thorough review of our article.

Financial support and sponsorship

G.L.T. reports a career development grant from the University of Zurich, Zurich, Switzerland (filling the gap program).

Conflicts of interest

There are no conflicts of interest.

REFERENCES AND RECOMMENDED READING

Papers of particular interest, published within the annual period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Liu GT, Schatz NJ, Galetta SL, *et al.* Persistent positive visual phenomena in migraine. *Neurology* 1995; 45:664–668.
2. Puledda F, Schankin C, Digre K, Goadsby PJ. Visual snow syndrome: what we know so far. *Curr Opin Neurol* 2018; 31:52–58.
- Excellent review on the history (including early reports), on the clinical characterization, differential diagnoses, and presumed pathophysiology of visual snow syndrome.
3. Rastogi RG, VanderPluym J, Lewis KS. Migrainous aura, visual snow, and 'alice in wonderland' syndrome in childhood. *Semin Pediatr Neurol* 2016; 23:14–17.
4. Simpson JC, Goadsby PJ, Prabhakar P. Positive persistent visual symptoms (visual snow) presenting as a migraine variant in a 12-year-old girl. *Pediatr Neurol* 2013; 49:361–363.
5. Schankin CJ, Goadsby PJ. Visual snow – persistent positive visual phenomenon distinct from migraine aura. *Curr Pain Headache Rep* 2015; 19:23.
6. Bou Ghannam A, Pelak VS. Visual snow: a potential cortical hyperexcitability syndrome. *Curr Treat Options Neurol* 2017; 19:9.
7. White OB, Clough M, McKendrick AM, Fielding J. Visual snow: visual misperception. *J Neuroophthalmol* 2018; 38:514–521.
- Very comprehensive review on visual snow syndrome, including demographic features and frequency of clinical findings, as well as discussion of individual symptoms in the context of pathophysiological considerations.
8. Literature commentary. *J Neuroophthalmol* 2019; 39:273–278.
9. Chen BS, Lance S, Lallu B, Anderson NE. Visual snow: not so benign. *J Clin Neurosci* 2019; 64:37–39.
10. Piquet AL, Khan M, Warner JEA, *et al.* Novel clinical features of glycine receptor antibody syndrome: a series of 17 cases. *Neurol Neuroimmunol Neuroinflamm* 2019; 6:e592.
11. Schankin CJ, Maniyar FH, Digre KB, Goadsby PJ. 'Visual snow' – a disorder distinct from persistent migraine aura. *Brain* 2014; 137(Pt 5):1419–1428.
12. Bessero AC, Plant GT. Should 'visual snow' and persistence of after-images be recognised as a new visual syndrome? *J Neurol Neurosurg Psychiatry* 2014; 85:1057–1058.
13. Lauschke JL, Plant GT, Fraser CL. Visual snow: a thalamocortical dysrhythmia of the visual pathway? *J Clin Neurosci* 2016; 28:123–127.
14. Schankin CJ, Maniyar FH, Sprenger T, *et al.* The relation between migraine, typical migraine aura and 'visual snow'. *Headache* 2014; 54:957–966.
15. Metzler AI, Robertson CE. Visual snow syndrome: proposed criteria, clinical implications, and pathophysiology. *Curr Neurol Neurosci Rep* 2018; 18:52.
- Case descriptions of primary and secondary visual snow, and review on visual snow syndrome suggesting a mild modification of diagnostic criteria. First article to mention the theory of stochastic resonance as a possible explanation for symptoms frequently associated with visual snow.
16. Kanai R, Paulus W, Walsh V. Transcranial alternating current stimulation (tACS) modulates cortical excitability as assessed by TMS-induced phosphene thresholds. *Clin Neurophysiol* 2010; 121:1551–1554.
17. Tegetmeyer H. Visual snow syndrome: symptoms and ophthalmological findings. *Klin Monbl Augenheilkd* 2017; 234:713–718.
18. Yildiz FG, Turkyilmaz U, Unal-Cevik I. The clinical characteristics and neurophysiological assessments of the occipital cortex in visual snow syndrome with or without migraine. *Headache* 2019; 59:484–494.
- Prospective clinical study investigating the habituation response by repetitive pattern reversal visual evoked potentials (VEP) and the phosphene threshold by transcranial magnetic stimulation. The study not only supports the hypothesis of occipital cortex hyperexcitability in visual snow but also provides a possible quantitative assessment tool in affected patients.
19. van Dongen RM, Waaijer LC, Onderwater GLJ, *et al.* Treatment effects and comorbid diseases in 58 patients with visual snow. *Neurology* 2019; 93:e398–e403.
- First systematic retrospective chart review on the benefit of pharmacological treatment in a large-case series of 58 patients.
20. Jeanmonod D, Schulman J, Ramirez R, *et al.* Neuropsychiatric thalamocortical dysrhythmia: surgical implications. *Neurosurg Clin N Am* 2003; 14:251–265.
21. Llinas RR, Ribary U, Jeanmonod D, *et al.* Thalamocortical dysrhythmia: a neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci U S A* 1999; 96:15222–15227.
22. Proske JH, Jeanmonod D, Verschure PFMJ. A computational model of thalamocortical dysrhythmia. *Eur J Neurosci* 2011; 33:1281–1290.
23. Chen WT, Lin YY, Fuh JL, *et al.* Sustained visual cortex hyperexcitability in migraine with persistent visual aura. *Brain* 2011; 134(Pt 8):2387–2395.
24. McKendrick AM, Chan YM, Tien M, *et al.* Behavioral measures of cortical hyperexcitability assessed in people who experience visual snow. *Neurology* 2017; 88:1243–1249.
25. Eren O, Rauschel V, Ruscheweyh R, *et al.* Evidence of dysfunction in the visual association cortex in visual snow syndrome. *Ann Neurol* 2018; 84:946–949.
- Clinical study using VEP and supporting the hypothesis of visual snow being a disorder involving the extrastriate visual cortex. The VEP findings attributed to the fusiform gyrus might be an electrophysiological correlate to the focal hypermetabolism of the adjacent lingual gyrus, which was demonstrated in a previous study on [(18) F]-2-fluoro-2-deoxy-D-glucose PET in patients with visual snow.
26. Di Russo F, Martinez A, Sereno MI, *et al.* Cortical sources of the early components of the visual evoked potential. *Hum Brain Mapp* 2002; 15:95–111.
27. Luna S, Lai D, Harris A. Antagonistic relationship between VEP potentiation and gamma power in visual snow syndrome. *Headache* 2018; 58:138–144.
28. Unal-Cevik I, Yildiz FG. Visual snow in migraine with aura: further characterization by brain imaging, electrophysiology, and treatment – case report. *Headache* 2015; 55:1436–1441.
29. Orland PM, Uglem M, Hagen K, *et al.* Visual evoked potentials in migraine: is the 'neurophysiological hallmark' concept still valid? *Clin Neurophysiol* 2016; 127:810–816.
30. Jager HR, Giffin NJ, Goadsby PJ. Diffusion- and perfusion-weighted MR imaging in persistent migrainous visual disturbances. *Cephalalgia* 2005; 25:323–332.
31. Schankin CJ, Viana M, Goadsby PJ. Persistent and repetitive visual disturbances in migraine: a review. *Headache* 2017; 57:1–16.
32. Trevino M, De la Torre-Valdovinos B, Manjarrez E. Noise improves visual motion discrimination via a stochastic resonance-like phenomenon. *Front Hum Neurosci* 2016; 10:572.
33. Manjarrez E, Mendez I, Martinez L, *et al.* Effects of auditory noise on the psychophysical detection of visual signals: cross-modal stochastic resonance. *Neurosci Lett* 2007; 415:231–236.
34. Wang YF, Fuh JL, Chen WT, *et al.* The visual aura rating scale as an outcome predictor for persistent visual aura without infarction. *Cephalalgia* 2008; 28:1298–1304.
35. Hampson M, Stoica T, Saksa J, *et al.* Real-time fMRI biofeedback targeting the orbitofrontal cortex for contamination anxiety. *J Vis Exp* 2012; e3535 10.3791/3535, doi: 10.3791/3535.
36. Kohler CA, Carvalho AF, Alves GS, *et al.* Autobiographical memory disturbances in depression: a novel therapeutic target? *Neural Plast* 2015; 2015:759139.
37. Lang PJ, McTeague LM, Bradley MM. Pathological anxiety and function/dysfunction in the brain's fear/defense circuitry. *Restor Neurol Neurosci* 2014; 32:63–77.
38. Mehler DMA, Sokunbi MO, Habes I, *et al.* Targeting the affective brain-a randomized controlled trial of real-time fMRI neurofeedback in patients with depression. *Neuropsychopharmacology* 2018; 43:2578–2585.
39. Yuan H, Young KD, Phillips R, *et al.* Resting-state functional connectivity modulation and sustained changes after real-time functional magnetic resonance imaging neurofeedback training in depression. *Brain Connect* 2014; 4:690–701.
40. Zhao Z, Yao S, Li K, *et al.* Real-time functional connectivity-informed neurofeedback of amygdala-frontal pathways reduces anxiety. *Psychother Psychosom* 2019; 88:5–15.
41. Zotev V, Yuan H, Misaki M, *et al.* Correlation between amygdala BOLD activity and frontal EEG asymmetry during real-time fMRI neurofeedback training in patients with depression. *Neuroimage Clin* 2016; 11:224–238.